arXiv:1405.5659v1 [math.AP] 22 May 2014

An elementary proof of asymptotic behavior of solutions of $u'' = Vu$

Giorgio Metafune* and Motohiro Sobajima†

Abstract. We provide an elementary proof of the asymptotic behavior of solutions of second order differential equations.

Mathematics Subject Classification (2010): 34E10.

Key words and phrases: Elementary proof, ordinary differential equations, asymptotic behavior.

1 Introduction

The asymptotic behaviour of the solutions of the ordinary differential equation

$$u''(x) = V(x)u(x), \qquad x \in (0,\infty) \tag{1.1}$$

is an important tool in various fields of mathematics and mathematical physics, in particular when special functions are involved. It can be found in [**3**, Section 6.2] and partially in [**1**, Chapter 10] and in [**2**, Chapter IV] when $V(x) = f(x) + g(x)$

$$u''(x) = \big(f(x) + g(x)\big)u(x), \qquad x \in (0,\infty) \tag{1.2}$$

assuming

$$\psi_{f,g} := |f|^{-\frac{1}{4}} \left(-\frac{d^2}{dx^2} + g\right) |f|^{-\frac{1}{4}} \in L^1(0,\infty). \tag{1.3}$$

The proof is usually done treating first the cases $f = \pm 1$ and then reducing to them the general case, by the Liouville transformation. We follow the same approach but simplify the cases $f = \pm 1$ by using Gronwall's Lemma, instead of successive approximations. In order to keep the exposition at an elementary level, we avoid also Lebesgue integration and dominated convergence (which could shorten some proofs). We consider both the behavior at infinity and near isolated singularities and apply the results to Bessel functions. We also recall that the general case

$$u''(x) + g(x)u'(x) = V(x)u(x)$$

can be reduced to the form (1.1) (with another V) by writing $u = \frac{1}{2}(\exp \int g)v$.

*Dipartimento di Matematica "Ennio De Giorgi", Università del Salento, Via Per Arnesano, 73100, Lecce, Italy, E-mail: giorgio.metafune@unisalento.it

†Dipartimento di Matematica "Ennio De Giorgi", Università del Salento, Via Per Arnesano, 73100, Lecce, Italy, E-mail: msobajima1984@gmail.com

2 Behavior near infinity in the simplest cases

First we consider the cases $f \equiv 1$ and $f \equiv -1$ and we prove the following results to which the general case reduces.

Proposition 2.1. *If $f = 1$, $g \in L^1(0,\infty)$, then there exist two solutions u_1 and u_2 of (1.2) such that, as $x \to \infty$,*

$$e^{-x}u_1(x) \to 1, \qquad e^{-x}u_1'(x) \to 1, \tag{2.1}$$

$$e^{x}u_2(x) \to 1, \qquad e^{x}u_2'(x) \to -1. \tag{2.2}$$

Proposition 2.2. *If $f = -1$, $g \in L^1(0,\infty)$, then there exist two solutions v_1 and v_2 of (1.2) such that, as $x \to \infty$,*

$$e^{-ix}u_1(x) \to 1, \qquad e^{-ix}u_1'(x) \to i, \tag{2.3}$$

$$e^{ix}u_2(x) \to 1, \qquad e^{ix}u_2'(x) \to -i. \tag{2.4}$$

By variation of parameters, every solution of (1.2) can be written as

$$u(x) = c_1 e^{\zeta x} + c_2 e^{-\zeta x} + \frac{1}{2\zeta}\int_a^x (e^{\zeta(x-s)} - e^{-\zeta(x-s)})g(s)u(s)\,ds, \quad x \in [a,\infty), \tag{2.5}$$

with $c_1, c_2 \in \mathbb{C}$, $\zeta = 1, i, -i$ and $a > 0$. In the following Lemma we choose $c_1 = 1, c_2 = 0$ to construct a solution which behaves like $e^{\zeta x}$ as $x \to \infty$, $\zeta = 1, i, -i$.

Lemma 2.3. *Let $\zeta \in \{1, i, -i\}$, $a > 0$ and $g \in L^1(a,\infty)$. If $u \in C^2([a,\infty))$ satisfies*

$$u(x) = e^{\zeta x} + \frac{1}{2\zeta}\int_a^x (e^{\zeta(x-s)} - e^{-\zeta(x-s)})g(s)u(s)\,ds, \qquad x \in [a,\infty),$$

then $z(x) := e^{-\zeta x}u(x)$ satisfies

$$|z(x)| \le e^{\int_a^x |g(r)|\,dr}, \qquad x \in [a,\infty) \tag{2.6}$$

$$\|zg\|_{L^1(a,\infty)} \le e^{\|g\|_{L^1(a,\infty)}} - 1. \tag{2.7}$$

Proof. Note that

$$z(x) = 1 + \frac{1}{2\zeta}\int_a^x (1 - e^{-2\zeta(x-s)})g(s)z(s)\,ds, \quad x \in [a,\infty).$$

Since $|1 - e^{-2\zeta(x-s)}| \le 2$ for $s \le x$, we see that for $x \ge a$,

$$|z(x)| \le 1 + \left|\frac{1}{2\zeta}\int_a^x (1 - e^{-2\zeta(x-s)})g(s)z(s)\,ds\right| \le 1 + \int_a^x |g(s)|\,|z(s)|\,ds.$$

Thus Gronwall's lemma implies (2.6), in particular z is bounded on $[a,\infty)$ and then $zg \in L^1(a,\infty)$. Moreover we have

$$\|zg\|_{L^1(a,\infty)} \le \int_a^\infty |g(s)|\,e^{\int_a^s |g(r)|\,dr}\,ds = e^{\|g\|_{L^1(a,\infty)}} - 1.$$

□

Proof of Proposition 2.1. Let $a > 0$ such that $\|g\|_{L^1(a,\infty)} < \log 2$ and let u be in Lemma 2.3 with $\zeta = 1$. Then u is one solution of (1.2) with $f = 1$. Set $z(x) = e^{-x}u(x)$. Then noting that as $x \to \infty$,

$$\begin{aligned}\left|\int_a^x e^{-2(x-s)}g(s)z(s)\,ds\right| &\le \int_a^{\frac{a+x}{2}} e^{-2(x-s)}|g(s)z(s)|\,ds + \int_{\frac{a+x}{2}}^x |g(s)z(s)|\,ds \\ &\le e^{-x+a}\|gz\|_{L^1(a,\infty)} + \|gz\|_{L^1(\frac{a+x}{2},\infty)} \to 0,\end{aligned}$$

we see that z satisfies

$$\begin{aligned} z(x) &\to z_\infty := 1 + \int_a^\infty g(s)z(s)\,ds \quad \text{as } x \to \infty, \\ z'(x) &= \int_a^x e^{-2(x-s)}g(s)z(s)\,ds \to 0 \quad \text{as } x \to \infty. \end{aligned}$$

By (2.7), we deduce that $\|zg\|_{L^1(a,\infty)} < 1$. Therefore $|z_\infty - 1| \le \|zg\|_{L^1(a,\infty)} < 1$ and hence $z_\infty \neq 0$. The function $u_1(x) := z_\infty^{-1}e^x z(x)$ satisfies (2.1). Moreover, since u_1^{-2} is integrable near ∞, another solution of (1.2) is given by

$$u_2(x) = 2u_1(x)\int_x^\infty \frac{1}{u_1(s)^2}\,ds. \tag{2.8}$$

Integrating by parts we deduce that, as $x \to \infty$,

$$\begin{aligned} e^x u_2(x) &= 2z_\infty e^{2x}z(x)\int_x^\infty \frac{1}{e^{2s}[z(s)]^2}\,ds \\ &= z_\infty e^{2x}z(x)\left(-\left[\frac{1}{e^{2s}[z(s)]^2}\right]_{s=x}^{s=\infty} - 2\int_x^\infty \frac{z'(s)}{e^{2s}[z(s)]^3}\,ds\right) \to 1 \end{aligned}$$

and

$$[e^x u_2(x)]' = 2z_\infty e^{2x}z'(x)\int_x^\infty \frac{1}{e^{2s}[z(s)]^2}\,ds + 2e^x u_2(x) - \frac{2z_\infty}{z(x)} \to 0.$$

□

Proof of Proposition 2.2. Let $a > 0$ such that $\|g\|_{L^1(a,\infty)} < \log 2$ and let $\tilde{u}_1$ and $\tilde{u}_2$ be as in Lemma 2.3 with $\zeta = i$ and with $\zeta = -i$, respectively. Noting that both $\tilde{u}_1$ and $\tilde{u}_2$ satisfy (1.2) with $f = -1$, and setting $z_1(x) = e^{-ix}\tilde{u}_1(x)$ and $z_2(x) = e^{ix}\tilde{u}_2(x)$, we have as $x \to \infty$

$$\begin{aligned} e^{2ix}\left(z_1(x) - 1 - \frac{1}{2i}\int_a^\infty g(s)z_1(s)\,ds\right) &\to \frac{1}{2i}\int_a^\infty e^{2is}g(s)z_1(s)\,ds, \\ e^{-2ix}\left(z_2(x) - 1 + \frac{1}{2i}\int_a^\infty g(s)z_2(s)\,ds\right) &\to -\frac{1}{2i}\int_a^\infty e^{-2is}g(s)z_2(s)\,ds \end{aligned}$$

and

$$e^{2ix}z_1'(x) \to \int_a^\infty e^{2is}g(s)z_1(s)\,ds, \qquad e^{-2ix}z_2'(x) \to \int_a^\infty e^{-2is}g(s)z_2(s)\,ds.$$

It follows that $\tilde{u}_1 \approx \xi_1 e^{ix} + \xi_2 e^{-ix}$, $\tilde{u}_1' \approx i\xi_1 e^{ix} - i\xi_2 e^{-ix}$ and $\tilde{u}_2 \approx \eta_1 e^{ix} + \eta_2 e^{-ix}$, $\tilde{u}_2' \approx i\eta_1 e^{ix} - i\eta_2 e^{-ix}$ as $x \to \infty$ where

$$\xi_1 = 1 + \frac{1}{2i}\int_a^\infty g(s)z_1(s)\,ds, \qquad \xi_2 = -\frac{1}{2i}\int_a^\infty e^{2is}g(s)z_1(s)\,ds,$$

and similarly for η_1, η_2. From (2.7) we see that $|\xi_1| > 1/2$, $|\xi_2| < 1/2$, $|\eta_1| < 1/2$ and $|\eta_2| > 1/2$ and hence $|\xi_1\eta_2 - \xi_2\eta_1| > 0$ and $\tilde{u}_1$ and $\tilde{u}_2$ are linearly independent. Therefore we can construct solutions u_1 and u_2 which satisfy (2.3) and (2.4), respectively. □

We consider now the case $f = 0$, assuming extra conditions on g.

Proposition 2.4. *Assume that $xg \in L^1(0,\infty)$. Then there exist two solutions u_1 and u_2 of*

$$u''(x) = g(x)u(x) \tag{2.9}$$

such that

$$\begin{aligned} x^{-1}u_1(x) &\to 1, \qquad & u_1'(x) &\to 1, \\ u_2(x) &\to 1, \qquad & xu_2'(x) &\to 0 \end{aligned}$$

as $x \to \infty$, respectively.

Proof. Set $u(x) := xz(x)$. Then $z'' + (2/x)z' = gz$ and, assuming $z'(a) = 0$ we obtain

$$z'(x) = x^{-2}\int_a^x s^2 g(s)z(s)\,ds. \tag{2.10}$$

Then assuming $z(a) = 1$

$$\begin{aligned} |z(x) - 1| &\le \int_b^x t^{-2}\left(\int_a^t s^2|g(s)z(s)|\,ds\right)\,dt \\ &= \int_a^x \left(\int_s^x t^{-2}\,dt\right) s^2|g(s)z(s)|\,ds \le \int_a^x s|g(s)z(s)|\,ds. \end{aligned} \tag{2.11}$$

Gronwall's lemma yields

$$|z(x)| \le e^{\int_a^x s|g(s)|\,ds}$$

hence z is bounded and $z' \in L^1(a,\infty)$ by (2.10). As in the proof of Proposition 2.1, $z(x) \to z_\infty \neq 0$ if a is sufficiently large. Moreover, since as $x \to \infty$,

$$|xz'(x)| \le \sqrt{\frac{a}{x}}\int_a^{\sqrt{ax}} s|g(s)z(s)|\,ds + \int_{\sqrt{ax}}^x s|g(s)z(s)|\,ds \to 0,$$

$u_1(x) := z_\infty^{-1}xz(x)$ satisfies the statement. Another solution u_2 of (1.2) is given by

$$u_2(x) := u_1(x)\int_x^\infty \frac{1}{u_1(s)^2}\,ds.$$

As in the proof of Proposition 3.1 we can verify that u_2 satisfies $u_2(x) \to 1$ and $xu_2'(x) \to 0$ as $x \to \infty$. □

Observe the integrability condition for xg near ∞ is necessary. In fact, if $g(x) = cx^{-2}$ the above equation has solutions x^α if $\alpha^2 - \alpha = c$.

3 Behavior near infinity in the general case

We recall that the function $\psi_{f,g}$ is defined in (1.3) and set $v_j(x) = |f|^{1/4}u_j(x)$, $j = 1, 2$ if u_1, u_2 are solutions of (1.2). The hypothesis $|f|^{1/2}$ not summable near ∞ guarantees that the Liouville transformation Φ of Lemma 3.3 maps (a, ∞) onto $(0, \infty)$, so that the results of the previous section apply. When it is not satisfied Φ maps (a, ∞) onto a bounded interval $(0, b)$ and the behavior of the solutions of (3.5) near b is more elementary (in some cases one can use Proposition 2.4).

Proposition 3.1. *Assume that $f(x) > 0$ in (a, ∞), $|f|^{1/2} \notin L^1(a, \infty)$ and $\psi_{f,g} \in L^1(a, \infty)$. Then there exist two solutions u_1 and u_2 of (1.2) such that as $x \to \infty$*

$$e^{-\int_a^x |f(r)|^{1/2}dr} v_1(x) \to 1, \qquad |f(x)|^{-1/2} e^{-\int_a^x |f(r)|^{1/2}dr} v_1'(x) \to 1, \tag{3.1}$$

$$e^{\int_a^x |f(r)|^{1/2}dr} v_2(x) \to 1, \qquad |f(x)|^{-1/2} e^{\int_a^x |f(r)|^{1/2}dr} v_2'(x) \to -1. \tag{3.2}$$

Proposition 3.2. *Assume that $f(x) < 0$ in (a, ∞), $|f|^{1/2} \notin L^1(a, \infty)$ and $\psi_{f,g} \in L^1(a, \infty)$. Then there exists two solutions u_1 and u_2 of (1.2) such that asx $\to \infty$*

$$e^{-i\int_a^x |f(r)|^{1/2}dr} v_1(x) \to 1, \qquad |f(x)|^{-1/2} e^{-i\int_a^x |f(r)|^{1/2}dr} v_1'(x) \to i, \tag{3.3}$$

$$e^{i\int_a^x |f(r)|^{1/2}dr} v_2(x) \to 1, \qquad |f(x)|^{-1/2} e^{i\int_a^x |f(r)|^{1/2}dr} v_2'(x) \to -i. \tag{3.4}$$

The proof is based on the well-known Liouville transformation that we recall below.

Lemma 3.3. *Let $a > 0$ and assume that $f \in C^2([a, \infty))$ satisfies $|f(x)| > 0$, $|f|^{1/2} \notin L^1(a, \infty)$. Define $\Phi \in C^2([a, \infty))$ by*

$$\Phi(x) := \int_a^x |f(r)|^{1/2}\, dr, \quad x \in [a, \infty).$$

Then $\Phi^{-1} : [0, \infty) \to [a, \infty)$ and if u satisfies (1.2) the function

$$w(y) := |f(\Phi^{-1}(y))|^{1/4} u(\Phi^{-1}(y)), \quad y \in [0, \infty)$$

satisfies

$$w''(y) = \left(\frac{f(\Phi^{-1}(y))}{|f(\Phi^{-1}(y))|} + \frac{\psi_{f,g}(\Phi^{-1}(y))}{|f(\Phi^{-1}(y))|^{1/2}} \right) w(y). \tag{3.5}$$

Proof. Note that $\Phi'(x) = |f(x)|^{1/2}$ and $\frac{d(\Phi^{-1})}{dy}(y) = |f(\Phi^{-1}(y))|^{-1/2}$. Setting $w(y) =$

$|f(\Phi^{-1}(y))|^{1/4}u(\Phi^{-1}(y))$ (and using $\xi = \Phi^{-1}(y)$ for simplicity), we have

$$\begin{aligned}
w'(y) &= \frac{d}{dx}\left[|f|^{1/4}u\right](\xi)\frac{d(\Phi^{-1})}{dy}(y) \\
&= |f(\xi)|^{-1/4}u'(\xi) + \left[|f|^{-1/2}\frac{d}{dx}|f|^{1/4}\right](\xi)u(\xi) \\
&= \left[|f|^{-1/4}u' - \frac{d}{dx}(|f|^{-1/4})u\right](\xi), \\
w''(y) &= \frac{d}{dx}\left[|f|^{-1/4}u' - \frac{d}{dx}(|f|^{-1/4})u\right](\xi)\frac{d(\Phi^{-1})}{dy}(y) \\
&= |f(\xi)|^{-3/4}u''(\xi) - \left[|f|^{-1/2}\frac{d^2}{dx^2}|f|^{-1/4}\right](\xi)u(\xi) \\
&= |f(\xi)|^{-1}(f(\xi)+g(\xi))w(y) - \left[|f|^{-3/4}\frac{d^2}{dx^2}|f|^{-1/4}\right](\xi)w(y).
\end{aligned}$$

Thus we obtain (3.5). □

Proof of Propositions 3.1 and 3.2. It suffices to apply Propositions 2.1 and 2.2 to the respective cases $f > 0$ and $f < 0$. Set $h(y) = \psi_{f,g}(\Phi^{-1}(y))|f(\Phi^{-1}(y))|^{-1/2}$. Then

$$\int_0^b |h(y)|\,dy = \int_a^\infty |\psi_{f,g}(x)|\,dx.$$

Therefore Propositions 2.1 and 2.2 are applicable to $w'' = \pm w + hw$, respectively. Finally, using Lemma 3.3 and taking $u(x) = |f(x)|^{-1/4}w(\Phi(x))$, we obtain the respective assertions in Propositions 3.1 and 3.2. □

4 Behavior near interior singularities

If f and g have local singularities at x_0, then the behavior of solutions near x_0 is also considerable. For simplicity, we take $x_0 = 0$. The following propositions are meaningful when $|f|^{1/2}$ is not integrable near 0, in particular when $|f|^{1/2} = cx^{-1}$. We recall that $v_j(x) = |f(x)|^{1/4}u_j(x)$, $j = 1, 2$.

Proposition 4.1. *Assume that $f(x) > 0$ in $(0,\infty)$ and $\psi_{f,g} \in L^1(0,\infty)$. Then there exist two solutions u_1 and u_2 of (1.2) such that as $x \downarrow 0$*

$$\begin{aligned}
e^{-\int_x^1 |f(r)|^{1/2}dr}v_1(x) &\to 1, \qquad |f(x)|^{-1/2}e^{-\int_x^1 |f(r)|^{1/2}dr}v_1'(x) \to -1, \\
e^{\int_x^1 |f(r)|^{1/2}dr}v_2(x) &\to 1, \qquad |f(x)|^{-1/2}e^{\int_x^1 |f(r)|^{1/2}dr}v_2'(x) \to 1.
\end{aligned}$$

Proposition 4.2. *Assume that $f(x) < 0$ in $(0,\infty)$ and $\psi_{f,g} \in L^1(0,\infty)$. Then there exist two solutions u_1 and u_2 of (1.2) such that as $x \downarrow 0$*

$$\begin{aligned}
e^{-\int_x^1 |f(r)|^{1/2}dr}v_1(x) &\to 1, \qquad |f(x)|^{-1/2}e^{-\int_x^1 |f(r)|^{1/2}dr}v_1'(x) \to -i, \\
e^{\int_x^1 |f(r)|^{1/2}dr}v_2(x) &\to 1, \qquad |f(x)|^{-1/2}e^{\int_x^1 |f(r)|^{1/2}dr}v_2'(x) \to i.
\end{aligned}$$

Proof of Propositions 4.1 and 4.2. Setting $w(s) := su(s^{-1})$ we see that

$$\begin{aligned} w''(s) &= s^{-3}u''(s^{-1}) \\ &= s^{-3}(f(s^{-1}) + g(s^{-1}))u(s^{-1}) = s^{-4}(f(s^{-1}) + g(s^{-1}))w(s). \end{aligned}$$

Let $\tilde{f}(s) := s^{-4}f(s^{-1})$ and $\tilde{g}(s) := s^{-4}g(s^{-1})$. Noting that

$$\begin{aligned} \psi_{\tilde{f},\tilde{g}}(s) &= s|f(s^{-1})|^{-1/4}\left(-\frac{d^2}{ds^2} + s^{-4}g(s^{-1})\right)\left(s|f(s^{-1})|^{-1/4}\right) \\ &= s^{-2}|f(s^{-1})|^{-1/4}\left(-\frac{d^2}{dx^2}|f|^{-1/4} + g|f|^{-1/4}\right)(s^{-1}) \\ &= s^{-2}\psi_{f,g}(s^{-1}), \end{aligned}$$

we have $\psi_{\tilde{f},\tilde{g}} \in L^1((0,\infty))$, and hence Propositions 3.1 and 3.2 can be applied. Since

$$\int_1^s |\tilde{f}(r)|^{1/2}dr = \int_{1/s}^1 |f(t)|^{1/2}dt,$$

we obtain the respective assertions in Propositions 4.1 and 4.2. □

5 Examples from special functions

Some examples illustrate the application of the results of the previous sections.

Example 1 (Modified Bessel functions)**.** We consider the modified Bessel equation of order ν

$$u'' + \frac{u'}{r} - \left(1 + \frac{\nu^2}{r^2}\right)u = 0, \tag{5.1}$$

All solutions of (5.1) can be written through the modified Bessel functions I_ν and K_ν. Both I_ν and K_ν are positive, I_ν is monotone increasing and K_ν is monotone decreasing (see e.g., [**3**, Theorem 7.8.1]). Proposition 2.1 and Proposition 4.1 give the precise behavior of I_ν and K_ν near ∞ and near 0, respectively. In fact, (5.1) can be written as

$$(\sqrt{r}u)'' = \left(1 + \frac{4\nu^2 - 1}{4r^2}\right)(\sqrt{r}u). \tag{5.2}$$

Since $1/r^2$ is integrable near ∞, choosing $f = 1$ and $g = \frac{4\nu^2-1}{4r^2}$, we see from Proposition 2.1 that

$$\sqrt{r}e^{-r}I_\nu(r) \to c_1 \neq 0 \quad \text{and} \quad \sqrt{r}e^{r}K_\nu(r) \to c_2 \neq 0 \qquad \text{as } r \to \infty.$$

Moreover, if $\nu \neq 0$, then choosing $f(r) = \frac{\nu^2}{r^2}$ and $g(r) = 1 - \frac{1}{4r^2}$, that is, $\psi_{f,g}(r) = r/\nu$, from Proposition 4.1 we have

$$r^{-\nu}I_\nu(r) \to c_3 \neq 0 \quad \text{and} \quad r^{\nu}K_\nu(r) \to c_4 \neq 0 \qquad \text{as } r \downarrow 0.$$

If $\nu = 0$, then putting $w(s) = u(e^{-s})$ we obtain

$$w''(s) = e^{-2s}w(s), \qquad s \in \mathbb{R}.$$

Therefore using Proposition 2.4 with $\tilde{g}(s) = e^{-2s}$ and taking $u(x) = w(-\log x)$, we have

$$I_0(r) \to c_5 \neq 0 \quad \text{and} \quad |\log r|^{-1} K_0(r) \to c_6 \neq 0 \qquad \text{as } r \downarrow 0.$$

Example 2 (Fundamental solution of $\lambda - \Delta$)**.** For $n \geq 3$, $\lambda \geq 0$ the fundamental solution v_λ of $\lambda - \Delta$ can be computed by integrating the heat kernel:

$$v_\lambda(r) = \int_0^\infty \frac{1}{(4\pi t)^{n/2}} e^{-\lambda t - \frac{r^2}{4t}}\, dt,$$

where $r = |x|$. Clearly $v_\lambda(r) \leq v_0(r) = cr^{2-n}$, $v_\lambda(r) \to 0$ as $r \to \infty$. The function $v = v_\lambda$ satisfies

$$v'' + \frac{n-1}{r} v' = \lambda v$$

or, setting $v = r^{(1-n)/2} w$,

$$w'' = \left(\lambda + \frac{n^2 - 1}{4r^2}\right) w.$$

Proceeding as in the example above we see that $r^{2-n} v(r) \to c_1 \neq 0$ as $r \to 0$ and $r^{(n-1)/2} e^{\sqrt{\lambda} r} v(r) \to c_2 \neq 0$ as $r \to \infty$.

Example 3 (Bessel functions)**.** Next we consider the Bessel equation of order ν

$$u'' + \frac{u'}{r} + \left(1 - \frac{\nu^2}{r^2}\right) u = 0, \tag{5.3}$$

or equivalently,

$$(\sqrt{r}u)'' = \left(-1 + \frac{4\nu^2 - 1}{4r^2}\right) (\sqrt{r}u).$$

All solutions of (5.3) can be written through the Bessel functions J_ν and Y_ν. As in Example 1, from Propositions 4.1 (for $\nu > 0$) and 2.4 (for $\nu = 0$) we obtain the behavior of J_ν and Y_ν near 0

$$r^{-\nu} J_\nu(r) \to c_1 \neq 0, \quad \text{and} \quad r^\nu Y_\nu(r) \to c_2 \neq 0 \qquad \text{as } r \downarrow 0$$

and if $\nu = 0$,

$$|\log r| J_0(r) \to c_3 \neq 0, \quad \text{and} \quad Y_0(r) \to c_4 \neq 0 \qquad \text{as } r \downarrow 0.$$

In view of Proposition 2.2 the behavior of J_ν and Y_μ near ∞ is given by

$$|\sqrt{r} J_\nu(r) - c_5 \cos(r + \theta_1)| \to 0, \quad \text{and} \quad |\sqrt{r} Y_\nu(r) - c_6 \cos(r + \theta_2)| \to 0,$$

as $r \to \infty$, where $c_5 \neq 0$, $c_6 \neq 0$ and $\theta_1, \theta_2 \in [0, \pi)$ satisfy $\theta_1 \neq \theta_2$.

References

[1] R. Beals, R. Wong, "Special functions," (A graduate text) Cambridge Studies in Advanced Mathematics **126**. Cambridge University Press, Cambridge, 2010.

[2] A. Erdely, "Asymptotic expansions," Dover 1956.

[3] F.W.J. Olver, "Asymptotics and special functions," Computer Science and Applied Mathematics, Academic Press, New York-London, 1974.